

## The Remainder Theorem and Synthetic Substitution/Division

© 2007-2008 Stephen Kent Stephenson. All rights reserved.

sks23@cornell.edu. Revised 3/5/08, 06:33.

Let  $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$  be a polynomial of degree  $n$ .

If  $P(x)$  is divided by  $x - b$ , then

$$\frac{P(x)}{x - b} = Q(x) + \frac{R}{x - b},$$

where  $Q(x)$  is the quotient, a polynomial of degree  $n - 1$ , and  $R$  is the remainder, a constant. Multiplying both sides by  $x - b$  gives

$$P(x) = (x - b)Q(x) + R.$$

If the polynomial is evaluated at  $x = b$ , then

$$P(b) = (b - b)Q(b) + R = 0 \cdot Q(b) + R = R, \text{ so}$$

$$P(x) = (x - b)Q(x) + P(b). \text{ This is } \mathbf{\textit{The Remainder Theorem}}.$$

**So, to find  $P(b)$  you can divide  $P(x)$  by  $x - b$  and the remainder will be  $P(b)$ .**

**If  $P(b) = 0$ , then  $x - b$  is a factor of  $P(x)$ .**

Of course you can always directly substitute  $b$  for  $x$  in the polynomial, like this:

$$P(b) = a_n b^n + a_{n-1} b^{n-1} + \dots + a_1 b + a_0,$$

$$P(b) = (((a_n)b + a_{n-1})b + \dots + a_1)b + a_0, \text{ which is called } \mathbf{\textit{The Nested Form}}.$$

The Nested Form represents a procedure that can be tabularized:

	$x^n$	$x^{n-1}$	$\dots$	$x^1$	$1$
$b$	$a_n$	$a_{n-1}$	$\dots$	$a_1$	$a_0$
(+)	$(a_n)b$	$\dots$	$(((a_n)b + a_{n-1})b + \dots + a_2)b$	$(((a_n)b + a_{n-1})b + \dots + a_2)b + a_1)b$	$(((a_n)b + a_{n-1})b + \dots + a_2)b + a_1)b$
	$a_n$	$(a_n)b + a_{n-1}$	$\dots$	$(((a_n)b + a_{n-1})b + \dots + a_2)b + a_1$	$P(b)$

This is known as *Synthetic Substitution*. For a concrete example, let  $f(x) = 2x^3 - 5x^2 + 7$ , then  $f(3)$  is found using Synthetic Substitution as follows in the figure at the left.

	$x^3$	$x^2$	$x^1$	$1$	
$3$	2	-5	0	7	$\leftarrow f(x)$
(+)		6	3	9	$\leftarrow bQ(x)$
	2	1	3	$16 = f(3)$	

  

	$x^3$	$x^2$	$x^1$	$1$	
$3$	2	-5	0	7	$\leftarrow f(x)$
(+)		6	3	9	$\leftarrow bQ(x)$
	2	1	3	$16 = f(3)$	

The Remainder Theorem can be used to understand what else is being shown in *Synthetic Substitution*. If  $P(x) = (x - b)Q(x) + P(b)$ , then  $P(x) + bQ(x) = xQ(x) + P(b)$ . So *Synthetic Substitution* also finds the quotient  $Q(x)$  of a division of  $P(x)$  by  $x - b$  (so *Synthetic Substitution* is also known as *Synthetic Division*), as annotated for our concrete example in the figure above right.